


Sexually Transmitted Disease Profile

Wahkiakum County 2004



Summary

This report describes the sexually transmitted disease burden in Wahkiakum County. Primary emphasis is placed on chlamydia and gonorrhea since they are the most frequently reported STDs in Washington State. The 2004 incidence rates by age and sex for gonorrhea and chlamydia are presented. The report concludes with a presentation of which providers in your county reported STDs.

Contents

County STD Trends

Table 1: Washington State Reportable Sexually Transmitted Diseases, 2004	1
--	---

Chlamydia

Figure 1: Chlamydia Incidence Rates, by Age and Gender, 2004.....	2
Table 2: Chlamydia Asymptomatic Infections, 2004	2

Gonorrhea

Figure 3: Gonorrhea Incidence Rates, by Age and Gender, 2004	4
Table 4: Gonorrhea Symptomatic Infections, 2004.....	4

Conclusion

Table 3: Reported Cases of Gonorrhea and Chlamydia by Provider Type, 2004	5
---	---

Appendix

A. Data Sources, Analyses, and Limitations	6
--	---

Wahkiakum County STD Disease Trends

Table 1: **Washington State** Reportable Sexually Transmitted Diseases, Wahkiakum County, 2004

Disease	2003 Wahkiakum County Cases	2004 Wahkiakum County Cases	2004 Wahkiakum County Rate ^λ (per 100,000)	2004 Washington State Rate ^λ (per 100,000)
Chlamydia	3	3	*	286
Gonorrhea	0	1	*	46
Early Syphilis	0	0	-	3.3
Congenital Syphilis	0	0	-	0.0 (live births)
Late/Late Latent Syphilis	0	0	-	2.2
Herpes (initial infection)	1	1	*	35
GI/LGV/Chancroid**	0	0	-	0.0
HIV cases**	0	0		
AIDS cases**	0	0		
TOTAL (excluding HIV/AIDS cases)	4	5	132	372

^λ Denominator estimates for the calculation of incidence rates from Washington State Adjusted Population Estimates, OFM, February 2004.

* Rates cannot be calculated for years with fewer than five cases.

** See Appendix A for explanation of disease acronyms.

2004 compared to 2003:

- Chlamydia had no change in reported cases (3 vs. 3).
- Initial infection herpes had a no change in reported cases (1 vs. 1).
- In 2004 - one (1) case of gonorrhea was reported. Zero (0) cases of gonorrhea were reported in 2003.

Complete information on the HIV/AIDS epidemic in Washington can be found in Washington State HIV/AIDS Surveillance Report, Washington State Department of Health, Office of IDRH Assessment Unit.

Chlamydia

Wahkiakum County Chlamydia rates by age cannot be calculated because all age groups have less than five cases in 2004.

Figure 1: **Chlamydia** Incidence Rates by Age and Gender, Wahkiakum County, 2004.

	0-9	10-14	15-19	20-24	25-29	30-34	35-39	40+
Female Rate	0	0	0	0	*	0	0	0
Male Rate	0	0	0	0	0	*	0	0
Female Cases	0	0	0	0	2	0	0	0
Male Cases	0	0	0	0	0	1	0	0

^λ Denominator estimates for the calculation of incidence rates from Washington State Adjusted Population Estimates, OFM, February 2004. Incidence rates rounded to the nearest whole number.

* Rates cannot be calculated for ages with fewer than five cases.

Women are preferentially screened for chlamydia. Because active case-finding is preferentially limited to women, the incidence of chlamydia in men may be under-reported by comparison. Caution should be used in interpreting comparisons of chlamydia rates between genders.

The 2002 STD Treatment Guidelines from CDC recommends that all women diagnosed with chlamydia be re-screened three to four months after treatment. This was suggested because of the high prevalence of chlamydia found in women diagnosed with the disease in the preceding months, presumably as a result of re-infection.

Asymptomatic Infection

STD infections often lack any signs and symptoms. Routine screening and treatment is essential to prevent serious complications that may not appear until long after infection. Screening all sexually active adolescents (19 years and younger) during sports physicals and routine office visits should be done even if symptoms are not present. Screening women and men aged 20-25 is also suggested, particularly those who have new or multiple sex partners. Women who are pregnant, have sex partners infected with chlamydia, have mucopurulent cervicitis or are planning an IUD insertion should also be screened. Careful interviewing and treatment of all partners is important.

Table 2: Reported Cases of **Chlamydia** by Diagnostic Category, Wahkiakum County, 2004.

Diagnosis	Private		Public		Total		Total Cases
	Male	Female	Male	Female	Male	Female	
Asymptomatic	0	0	0	0	0	0	0
Symptomatic-Uncomplicated	1	2	0	0	1	2	3
Pelvic Inflammatory Disease	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0
Unknown	0	0	0	0	0	0	0
TOTAL	1	2	0	0	1	2	3

Gonorrhea

There was one (1) case of gonorrhea reported in Wahkiakum County in 2004. Wahkiakum County rates for gonorrhea by age groups cannot be calculated because all age groups have less than five cases reported in 2004.

In Washington State the reported rate of gonorrhea incidence in 2004 was 46/100,000, a slight increase from the 2003 rate. Statewide, the greatest incidence of disease among females was among 15-19 year olds (198/100,000), while for males the burden of disease is distributed more evenly among those older. Males had a higher gonorrhea rate (51/100,000) than females (40/100,000). A major factor contributing to the distribution of gonorrhea incidence in different age groups among men or women is a documented outbreak among men who have sex with men (MSM) whose median reported age was 30.

Findings from the Gonococcal Isolate Surveillance Project (GISP) in Seattle have indicated that Washington State is now an area with increased prevalence of quinolone-resistant *Neisseria gonorrhoeae* (QRNG). Based on these findings, the Washington State Department of Health recommends that health care providers in the state should no longer use fluoroquinolones (ciprofloxacin, levofloxacin and ofloxacin) as first line therapy for gonorrhea. The antibiotics of choice are ceftriaxone (Rocephin™) or cefpodoxime (Vantin™) accompanied by either azithromycin or doxycycline to treat possible coexisting chlamydial infection.

Because most gonorrhea infections cause symptoms and prompt individuals to seek medical care, reported cases are considered to be an accurate reflection of true disease incidence in the overall population. Providers in Washington State who reported gonorrhea cases in 2004 indicated that 80% of the men were symptomatic for gonorrhea; 43% of the women were symptomatic.

Conclusion

Table 3: Reported Cases of Chlamydia and Gonorrhea by Provider Type, Wahkiakum County, 2004

Provider Type	Chlamydia			Gonorrhea		
	No. of Providers	No. of Cases	Percent of Total Cases	No. of Providers	No. of Cases	Percent of Total Cases
Alcohol/Substance Abuse						
Blood Bank/Plasma Center						
Community Health Center						
Emergency Care (excl. hosp.)						
Family Planning						
Health Plan/HMOs						
HIV/AIDS						
Hospitals	2	2	67%			
Indian Health						
Jail/Correction/Detention						
Job Corps						
Migrant Health						
Military						
Neighborhood Health						
OB/GYN				1	1	100%
Other	1	1	33%			
Private Physicians						
Reproductive Health						
STD Clinics						
Student Health						
TOTAL	3	3	100%	1	1	100%

In Wahkiakum County, the hospitals providers reported the highest number of chlamydia cases. These providers reported 67% of the total. Other reported the second highest number of chlamydia cases (33%).

In 2004, 100% of chlamydia and 0% of gonorrhea cases indicating treatment date by health care providers in Wahkiakum County were reported within 30 days to the STD Washington State Morbidity System.

The **Healthy People 2010** national objectives for chlamydia incidence are:

Females aged 15-24 attending family planning clinics: 3%.

There are zero (0) Region X Infertility Prevention Project (IPP) family planning clinics in Wahkiakum County.

Females aged 15-24 attending STD clinics: 3%.

Males aged 15-24 attending STD clinics: 3%.

There are zero (0) Region X IPP STD/reproductive health clinics inWahkiakum County.

The **Healthy People 2010** national objective for gonorrhea incidence is 19 cases per 100,000.

Wahkiakum County rates cannot be calculated because less than five cases were reported in 2004.

Appendix A: Data Sources, Analyses and Limitations

Cases: The number of cases identified and submitted by providers to local health jurisdictions and forwarded to the Washington State Department of Health, Office of Infectious Disease and Reproductive Health, STD/TB Services.

Population: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM), February 2004.

Incidence Rates: Incidence rates are calculated as the number of new episodes of a disease (not persons) in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates should not be calculated for incident case totals fewer than five because the rates are unstable.

Data Reporting: Gonorrhea, chlamydia, syphilis, and herpes (initial infection) are reportable diseases to the local health jurisdictions and forwarded to the Department of Health. To be reported and included in surveillance data, disease definition must be met.

Disease Definitions:

- AIDS - Acquired Immunodeficiency Syndrome is the advanced stage of HIV-disease in humans and is characterized by severe suppression of immune response. Persons with AIDS are at risk for increased susceptibility to opportunistic infections, degradation of major organ systems and eventual death.
- Chancroid - a STD characterized by painful genital ulceration and inflammatory inguinal adenopathy caused by the bacterium *Haemophilus ducreyi*.
- Chlamydia- isolation of *Chlamydia trachomatis* from a clinical specimen by culture or non-culture methods that detect chlamydia antigen or genetic material.
- Gonorrhea - isolation of *Neisseria gonorrhoeae* from a clinical specimen by culture or non-culture methods or observation of Gram-negative intracellular diplococci in urethral or endocervical smears.
- Granuloma Inguinale (GI) - a slowly progressive ulcerative disease of the skin and lymphatics of the genital and perianal area.
- Herpes Simplex (initial infection only) - diagnostic criteria for reporting can be made through clinical observation of typical lesions and/or laboratory confirmation.
- HIV - Human Immunodeficiency Virus is a retrovirus causing HIV disease and AIDS in humans. This pathogen is transmitted from person to person through unprotected sexual contact, sharing of injection equipment and transfusion/transplantation with infected blood or tissue.
- Lymphogranuloma Venereum (LGV) - characterized by genital lesions, suppurative regional lymphadenopathy, or hemorrhagic proctitis, caused by the L1, L2 and L3 serovars of *Chlamydia trachomatis*.

- Syphilis - a complex sexual transmitted disease with a highly variable clinical course. See CDC guidelines for surveillance definition.

The diagnosing practitioner is responsible for providing the case information which includes patient demographics, source of diagnosis, limited clinical information including site of infection and treatment, and date of diagnosis.

Data Strengths: Sexually transmitted disease data may provide more timely information on behavioral trends in the community than diseases with similar modes of transmission particularly HIV/AIDS. There is a high level of participation in the STD surveillance system by private providers of STD services.

Data Limitations: Clinically diagnosed cases of STDs (without laboratory confirmation) may be missed through this surveillance system. Depending upon diagnosing practices, completeness of reporting may vary by source of health care.

Data Biases: Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. However, it is assumed that the number of cases that would fall into these categories is small and normally distributed, thus not significantly impacting the calculated STD rates.

Assumptions: It is assumed that the cases reported from year to year are independent of each other. One violation of this assumption could be if a person who has an STD one year is more likely to have an STD the following year. Also, repeat episodes of the same STD by the same person are not excluded from the numerator count; it is felt that these numbers are not large enough to significantly impact the calculated incidence rates. Finally, we have assumed that all rates follow a chi-square distribution.

Region X IPP screening criteria that are used at the 140 IPP clinics are as follows:

- Sexually active women 24 years and younger;
- Pregnant women;
- Women with mucopurulent cervicitis, cervical friability, or ectopy with inflammation or edema;
- Women with pelvic inflammatory disease (PID);
- Women planning to receive an intrauterine device;
- Women with a symptomatic sex partner;
- Women diagnosed with CT in the last 12 months; and
- Sex partners of persons with chlamydial infection.